

2016-7403
2017-02-28

***Serenade*[®] *Opti*[™]**
A Wettable Powder Biofungicide

COMMERCIAL

GUARANTEE: QST 713 strain of dried *Bacillus subtilis*, contains a minimum of 1.31×10^{10} CFU/g

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

POTENTIAL SENSITIZER

CAUTION – EYE IRRITANT

REGISTRATION NO. 31666 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 kg - bulk

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. S.E.
Calgary, Alberta T2C 3G3

Product Information: 1-888-283-6847
www.bayercropscience.ca

Date of Manufacture:

Made in Mexico

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and *Pest Control Product* Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May irritate eyes. May cause sensitization. Avoid contact with eyes, skin or clothing. Avoid inhaling/breathing dusts. Wear a long-sleeved shirt, long pants, shoes plus socks, water-proof gloves and NIOSH approved mist filtering mask or respirator when handling, mixing/loading or applying the product and during all clean-up/repair activities.

STORAGE AND DISPOSAL:

To prevent contamination, store this product away from food or feed.

STORAGE: Store in a dry area inaccessible to children. Store at room temperature for up to 6 years from the date of manufacture. Store in original container only. Keep container closed when not in use.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory requirements. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

ENVIRONMENTAL HAZARDS

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or

disposal of wash water.

To reduce the runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

GENERAL USE INFORMATION:

Serenade Opti is a broad spectrum, preventative product for the suppression of many plant diseases. Serenade Opti may be applied as a foliar spray alone, or in an alternating spray program with other registered crop protection products. When conditions are conducive to heavy disease pressure, use Serenade Opti in a rotational program with other registered fungicides. Serenade Opti may be applied with ground or aerial application equipment.

INTEGRATED PEST MANAGEMENT (IPM):

Integrate Serenade Opti into an overall disease and pest management strategy whenever fungicide use is necessary. Follow practices known to reduce disease development. Consult local agricultural authorities for specific IPM strategies developed for your crop(s) and location.

USE RATE DETERMINATION:

Carefully read and follow all label directions, use rates and restrictions. Serenade Opti should be applied prior to or in the early stages of disease development. Use maximum label rates and shortened spray intervals for conditions conducive to rapid disease development. For proper application, determine the number of hectares to be treated, the recommended label use rate and select appropriate water volumes to give good canopy penetration and coverage of plant parts to be protected. Prepare only the amount of spray solution required to treat the measured acreage. Accurate spray equipment calibration is essential prior to use.

PREHARVEST INTERVAL:

Serenade Opti can be applied up to and including the day of harvest.

APPLICATION INSTRUCTIONS – Agricultural Use:

GENERAL: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions. Where provinces/territories have more stringent regulations, they should be observed.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

GROUND: Thorough coverage is essential for optimum disease control. To achieve good coverage, use the appropriate spray pressure, liters per hectare, nozzles, nozzle spacing and ground speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

AIR: Use the application rate indicated for the appropriate crop in sufficient water to achieve thorough coverage.

MIXING INSTRUCTIONS:

MIXING: Serenade Opti must be diluted with water for spray applications and may be used in spray equipment commonly used for making ground applications. Partially fill the spray tank with clean water and begin agitation. Add the specified amount of Serenade Opti to the tank. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation continuously while spraying. Do not allow spray mixture to stand overnight or for prolonged periods.

Application Rates for Serenade Opti

Directions for Use: When applied as directed (see table below), Serenade Opti will suppress the indicated diseases. Under conditions of moderate to high disease pressures, use the higher rate and shorter application interval indicated. Apply in sufficient water carrier volume to ensure good coverage. Serenade Opti has a 0 day PHI, and can be applied up to, and including, the day of harvest.

Ground Application Only.

Crop Group 1 Root tuber vegetables: Garden beet, sugar beet, carrot, celeriac, chervil, chicory, ginseng, horseradish, turnip-rooted parsley, parsnip, potato, radish, Oriental radish, rutabaga, salsify, sweet potato, and turnip.

Disease	Rate (kilogram/ hectare)	Application Instructions
White mould (<i>Sclerotinia sclerotiorum</i>)	1.1 – 2.2	Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval.

Celery

Disease	Rate (kilogram/ hectare)	Application Instructions
Pink Rot (<i>Sclerotinia sclerotiorum</i>)	1.7 – 3.3	Make the first application approximately 8 weeks before harvest and repeat on a 14-day interval. Apply Serenade Opti as a directed spray in sufficient water to ensure thorough coverage of the base of the plants and the surrounding soil surface.

Potato

Disease	Rate (kilogram/ hectare)	Application Instructions
Early blight (<i>Alternaria solani</i>)	1.1 – 2.2	Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval.
Silver Scurf (<i>Helminthosporium solani</i>)	7 - 14 g/tonne	<p>Potatoes: For post-harvest application to aid in the control of silver scurf. Sanitation and other cultural practices should also be employed to aid in control and minimize the potential for disease.</p> <p><u>Conveyer Line Application:</u> Prepare the equivalent of 350 – 700 grams of Serenade Opti in 100 liters of water. Spray 2 liters of the Serenade Opti/water suspension per tonne of potatoes. Potatoes must rotate along the conveyer line into the storage area to ensure complete coverage. If needed, adjust rate of spray solution to ensure thorough coverage while maintaining recommended rate of Serenade Opti per tonne of potatoes.</p>

Radish, turnips and rutabaga

Disease	Rate (kilogram/ hectare)	Application Instructions
Downy mildew (<i>Peronospora parasitica</i>)	1.4	Downy mildew: Begin application when environmental conditions are conducive to disease development and repeat on 7-10 day intervals.

Sugar, Garden Beet

Disease	Rate (kilogram/ hectare)	Application Instructions
Cercospora Leaf Spot (<i>Cercospora beticola</i>)	0.5 -1.1	<p>Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval.</p> <p>Serenade Opti must be applied with a non-ionic surfactant at a rate of 0.25% v/v or at labeled rates.</p> <p>Under conditions of moderate to high disease pressures, use the higher rate and shorter application interval indicated.</p>

Crop Group 3 - Bulb Vegetables: Onions (all types), garlic (all types), leeks, shallots and chives

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis neck rot (<i>Botrytis allii</i>) Botrytis leaf blight (<i>Botrytis squamosa</i>)	1.7 – 2.5	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
Downy mildew (<i>Peronospora destructor</i>)	1.7 – 3.3	

Crop Group 4 - Leafy Vegetables (except brassica): Amaranth, arugula, Chinese celery, celtuce, chervil, corn salad, cress, dandelion, dock, endive, fennel, lettuce, orach, parsley, radicchio, and swiss chard.

Disease	Rate (kilogram/ hectare)	Application Instructions
Sclerotinia rot (<i>Sclerotinia minor</i> , <i>Sclerotinia sclerotiorum</i>)	0.6 – 1.7	Make the first application at planting (or immediately following planting but prior to crop emergence). Make a second application as a directed spray with multiple nozzles per seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on 10-14 day intervals if conditions for disease development persist.
Grey mould (<i>Botrytis cinerea</i>)		For suppression, begin applications soon after emergence or transplant and continue as necessary on a 7 to 10 day interval. When environmental conditions are conducive to rapid disease development, use Serenade Opti in a rotational program with other registered fungicides. Thorough coverage is essential.
Powdery mildew (<i>Erysiphe cichoracearum</i>)	1.7 – 3.3	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Lettuce

Disease	Rate (kilogram/ hectare)	Application Instructions
Downy mildew (<i>Bremia lactucae</i>)	1.7 – 3.3	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Spinach

Disease	Rate (kilogram/ hectare)	Application Instructions
White rust (<i>Albugo occidentalis</i>)	0.6 – 1.1	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Crop Group 5 - Brassica Leafy Vegetables : Broccoli, Chinese broccoli, Broccoli raab, Brussels sprouts, cabbage (all types), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens.

Disease	Rate (kilogram/ hectare)	Application Instructions
Sclerotinia rot (<i>Sclerotinia sclerotium</i>)	1.7 – 3.3	Make the first application at emergence, or immediately following transplanting. Repeat applications on 10-14 day intervals if conditions for disease development continue.
Downy mildew (<i>Peronospora parasitica</i>)	1.1 – 1.7	
Pin rot (<i>Alternaria / Xanthomonas Complex</i>)		

Crop Group 6 – legume Vegetables: Bean (*Lupinus spp.*, *Phaseolus spp.*, and *Vigna spp.*), chickpea, lentil, and pea (all types)

Disease	Rate (kilogram/ hectare)	Application Instructions
White mould (<i>Sclerotinia sclerotiorum</i>)	1.7 – 3.3	Make the first application at planting (or immediately following planting but prior to crop emergence). Make a second application as a directed spray with multiple nozzles per seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on 7-14 day intervals if conditions for disease development persist.
Botrytis pod rot (<i>Botrytis cinerea</i>)	1.7 – 3.3	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Soybeans

Disease	Rate (kilogram/ hectare)	Application Instructions
White mould (<i>Sclerotinia sclerotium</i>)	0.6 – 2.0	Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval.
Brown spot (<i>Septoria glycines</i>)	0.1 – 0.5	
Frog-eye leaf spot (<i>Cercospora sojina</i>)		

Asparagus

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis blight (<i>Botrytis cinerea</i>)	1.7 – 3.3	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Crop Group 8 - Fruiting Vegetables: Eggplant, ground cherry, okra, pepino, peppers (all varieties), tomatillo and tomatoes

Disease	Rate (kilogram/ hectare)	Application Instructions
Early blight (<i>Alternaria solani</i>)	2.5	Make the first application when plants are 6-10 cm high, or when conditions are conducive for disease development. Repeat applications on an interval of 5-7 days.
Botrytis grey mould (<i>Botrytis cinerea</i>)	1.7 – 3.3	

Tomato, Peppers

Disease	Rate (kilogram/ hectare)	Application Instructions
Powdery mildew (<i>Erysiphe orontii</i> , <i>Leveillula taurica</i>)	1.7 – 3.3	Begin application soon after emergence or transplant, and repeat on an interval of 7-10 days.
Bacterial spot (<i>Xanthomonas campestris</i>)	0.6 – 1.7	Begin application soon after emergence or transplant, when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval. When environmental conditions and plant stage are conducive to rapid disease development, use Serenade Opti in a rotational program with other registered bacteriacides.

Crop Group 9 - Cucurbit Vegetables: Canteloupe, Chinese waxgourd, cucumber, gherkin, edible gourd, melon, muskmelon, pumpkin, squash, and watermelon.

Disease	Rate (kilogram/ hectare)	Application Instructions
Powdery mildew (<i>Erysiphe cichoracearum</i> , <i>Sphaerotheca fuliginea</i>)	1.7 – 3.3	Begin application soon after emergence or transplant, when conditions are conducive for disease development. Repeat applications on an interval of 7-10 days.
Downy mildew (<i>Pseudoperonospora cubensis</i>)	0.6 – 1.7	Begin application soon after emergence or transplant, when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval. When environmental conditions and plant stage are conducive to rapid disease development, use Serenade Opti in a rotational program with other registered fungicides.

Crop Group 11 - Pome Fruit: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental pear, Quince, and non-bearing pome fruit trees

Disease	Rate (kilogram/ hectare)	Application Instructions
Fire blight (<i>Erwinia amylovora</i>)	1.1 – 1.7	Begin applications at 1-5% bloom and repeat on 7-10 days interval. During periods of rapid bloom development and frequent infection periods, spray intervals of 4-7 days may be required.
Scab (<i>Venturia inaequalis</i> ; <i>V. pirina</i>)	1.7 – 3.3	Begin application at green tip or when environmental conditions become favorable for primary scab development, and repeat on 7-10 day intervals.

Apples

Disease	Rate (kilogram/ hectare)	Application Instructions
Powdery Mildew (<i>Podosphaera leucotricha</i>)	1.7 – 3.3	Begin application at tight cluster, or sooner, if conditions are conducive to disease development. Repeat applications on a 7-10 day interval. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure.

Crop Group 12 – Stone Fruit: Apricot, cherry, nectarine, peach, plum, plumcot, and prune

Disease	Rate (kilogram/ hectare)	Application Instructions
Brown Rot (<i>Monilinia spp.</i>)	1.1 – 1.7	Begin application at early bloom and repeat as necessary through petal fall on a 7-day interval.

Crop Group 13 – Small Fruit & Berries: Bayberry; bearberry; blackberry; highbush and lowbush blueberry; chokecherry; currant; elderberry; gooseberry; huckleberry; lingonberry; loganberry, mulberry; pincherry; raspberry; salal; Saskatoon berry; sea buckthorn; wild raspberry

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis grey mould (<i>Botrytis cinerea</i>)	1.7 - 3.3	Begin applications at the first sign of disease or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
Bacterial blight (<i>Pseudomonas syringae</i>)	0.6 – 1.7	Apply before fall rains and again during dormancy before spring.

Crop Group 13-07 – Berry and Small Fruit: Blackberry or raspberry, highbush blueberry, elderberry or mulberry, grape, strawberry, and fuzzy kiwifruit

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis grey mould mould (<i>Botrytis cinerea</i>)	1.7 – 3.3	Begin application prior to disease development and repeat on 7-10 day intervals.

Grapes

Disease	Rate (kilogram/ hectare)	Application Instructions
Powdery mildew (<i>Erysiphe necator</i> syn. <i>Uncinula necator</i>)	1.7 – 3.3	Begin application when new shoots are 1-3 cm high. Make a repeat application when shoots are 6-10 cm high, and when shoots are 18-20 cm high, then at 7-10 day intervals thereafter.
Grey mold (<i>Botrytis cinerea</i>)	1.7 – 3.3	Make the first application at bloom, then repeat at bunch closure, at veraison and pre-harvest.
Sour rot* (a complex of various pathogens)		*Note that sour rot disease is caused by various yeast, fungal and bacterial pathogens, and that Serenade Opti has not been tested for control of all of these agents.

Highbush blueberry, lowbush blueberry

Disease	Rate (kilogram/ hectare)	Application Instructions
Mummy berry (<i>Monilinia vaccinii-corymbosi</i>)	2.0 -3.3	Begin applications at the bud break stage of development. Repeat as necessary on a 7-14 day interval.

Strawberry

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis gray mould (<i>Botrytis cinerea</i>)	1.7 – 3.3	Begin application at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.

Crop Group 19 – Herbs and Spices: Anise, balm, basil, borage, chamomile, caraway, catnip, celery seed, chervil, chive, coriander, costmary, dill, fennel, fenugreek, horehound, juniper berry, lavender, lemongrass, lovage, marigold, mustard (seed), nasturtium, parsley, rosemary, sage, savory, sweet bay, thyme, wintergreen, woodruff, and wormwood including transplants

Disease	Rate (kilogram/ hectare)	Application Instructions
Botrytis gray mould (<i>Botrytis cinerea</i>); White mould (<i>Sclerotinia sclerotiorum</i>)	1.7 – 3.3	Begin application when environmental conditions are conducive to disease development. Repeat as necessary on a 7- to 10-day interval.

Peanut

Disease	Rate (kilogram/ hectare)	Application Instructions
Leaf spot (<i>Cercospora arachidicola</i> , <i>Cercosporidium personatum</i>)	0.6 – 1.7	Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat as necessary on a 7-10 day interval.

Ground and Aerial application.

Crop Group 20A - Rapeseed Subgroup. Representative commodity: rapeseed (canola varieties only) borage, crambe, cuphea, echium, flax seed, gold of pleasure, hare's ear mustard, lesquerella, lunaria, meadowfoam, milkweed, mustard seed, oil radish, poppy seed, rapeseed, sweet rocket, cultivars, varieties and/or hybrids of these

Disease	Rate (kilogram/hectare)	Application Instructions
Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	0.3 -0.9	Begin application at 20%- 30% bloom. A second application may be made 7-10 days later, at approximately 50% bloom and prior to significant petal fall, if conditions for disease development remain favorable. Use higher rates in fields with a history of heavy disease pressure. Aerial Spray Volume: minimum 50 L/ha

Aerial Application Instructions

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where not rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices and/or electronic tracking equipment

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only when meteorological conditions are in compliance with local and/or provincial authorities.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear the personal protective equipment described in the PRECAUTIONS section of this label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

MINOR USES:

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (BELOW): The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Bayer CropScience Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Bayer CropScience Inc. itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below. Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Bayer CropScience Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

Crop	Disease	Rate (kilogram/ hectare)	Application Instructions
Hops	Downy mildew (<i>Pseudoperonospora humuli</i>) Partial suppression	3.3	Downy mildew: Begin application when environmental conditions are conducive to disease development and repeat on 7-10 day intervals. Use the shorter application intervals under conditions of moderate to high disease pressure.
Alfalfa	Blossom blight (<i>Botrytis cinerea</i>) Suppression	1.7 – 3.3	Begin application prior to disease development when environmental conditions are conducive to disease development. Use higher rate and shorter application intervals under conditions of moderate to high disease pressure. Repeat as necessary on a 7- to 10-day interval.

160601

Serenade Opti is a registered trademark of Bayer.
© Copyright Bayer

***Serenade*[®] *Opti*[™]**
A Wettable Powder Biofungicide

COMMERCIAL

GUARANTEE: QST 713 strain of dried *Bacillus subtilis*, contains a minimum of 1.31×10^{10} CFU/g

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

POTENTIAL SENSITIZER

CAUTION – EYE IRRITANT

REGISTRATION NO. 31666 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 kg - bulk

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. S.E.
Calgary, Alberta T2C 3G3

Product Information: 1-888-283-6847
www.bayercropscience.ca

Date of Manufacture:

Made in Mexico

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and *Pest Control Product* Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May irritate eyes. May cause sensitization. Avoid contact with eyes, skin or clothing. Avoid inhaling/breathing dusts. Wear a long-sleeved shirt, long pants, shoes plus socks, water-proof gloves and NIOSH approved mist filtering mask or respirator when handling, mixing/loading or applying the product and during all clean-up/repair activities.

ENVIRONMENTAL HAZARDS

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wash water.

To reduce the runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE AND DISPOSAL:

To prevent contamination, store this product away from food or feed.

STORAGE: Store in a dry area inaccessible to children. Store at room temperature for up to 6 years from the date of manufacture. Store in original container only. Keep container closed when not in use.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory requirements. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

Serenade Opti is a registered trademark of Bayer.