

28-SEPT-2009
2009-2505

AGRICULTURAL

NET CONTENTS: 200 g
600 g
1 kg

AGROX® CD
DUAL PURPOSE SEED TREATMENT POWDER
DANGER POISON

CORN, BEANS, SOYBEANS AND PEAS:

For the control of seed corn maggots and to supplement previous fungicidal treatment for seedling blight and seed rot.

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN
DO NOT APPLY BY AIR

GROUP	1B	INSECTICIDE
GROUP	M	FUNGICIDE

CAUTION: Exposure to captan may produce long-term health effects. To minimize exposure, follow directions outlined on this label.

Warning, contains the allergen soy .

GUARANTEE: Diazinon 15%

Captan 15%

REGISTRATION NO. 26957 PEST CONTROL PRODUCTS ACT

NORAC CONCEPTS INC.
P.O. Box/C.P. 31097
GUELPH, ON N1H 8K1
24 hour inquiry phone no.: 519-821-3633

J00/A-1

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Do not breath dust. Avoid spilling on skin or clothing. When treating or handling treated seed, work in a well-ventilated area, and wear a respirator, goggles, and rubber gloves. Do not use leather or cloth gloves. Wear clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before reuse. Wash hands, face and arms thoroughly after handling AGROX CD and before drinking, eating or smoking. Shower after use or at the end of the work day. Do not contaminate food, feed or any body of water. Label unplanted treated seed as follows: "Poisonous to humans and animals. Do not use for food or feed purposes. Do not sell to oil mills. This seed has been treated with captan and diazinon."

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca

FIRST AID

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If swallowed**, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Diazinon is a cholinesterase inhibitor. Atropine is antidotal. Symptoms of Poisoning: nausea, vomiting, hyper-irritability, convulsions and/or coma and other symptoms typical to organophosphate insecticide poisoning.

EMERGENCY TELEPHONE NUMBER

All hours, (519) 821-3633 **ONLY** for health and environmental information.

DISPOSAL

1. Thoroughly empty the contents of the container into the application device.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency.
Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

RESISTANCE-MANAGEMENT RECOMMENDATIONS: For resistance management, please note that AGROX CD contains both a Group 1B insecticide, and Group M fungicide. Any insect or fungal population may contain individuals naturally resistant to AGROX CD and other Group 1B or Group M insecticides/fungicides. A gradual or total loss of insect control may occur over time if the insecticide is used repeatedly in the same fields. Also, gradual or total loss of insect or fungal control may occur over time if this fungicide is used repeatedly in the same fields. **To delay insecticide/fungicide resistance:** Where possible, rotate the use of AGROX CD with different groups that control the same pathogen; monitor treated insect and fungal populations for resistance development; if resistance is suspected, report it to the company representative, switch to another insecticide/fungicide with a different target site of action if possible, or consult a certified crop advisor or local extension specialist for guidance.

USE THIS PRODUCT ONLY ON SEED PREVIOUSLY TREATED WITH CAPTAN OR THIRAM. DO NOT USE ON SEED ALREADY TREATED WITH AN INSECTICIDE (other than methoxychlor or malathion). Corn, Beans, Soybeans and Peas: For the control of seed corn maggots and to supplement previous fungicidal treatment for Seedling Blight and Seed Rot, apply the contents of one 200 g container to 100 kg (4 units) of seed at planting or near planting time (50 g per 25 kg of seed). When starting with an empty planter, place enough treated seed in the planter box to cover the bottom. Treat this seed in a separate container that later can be destroyed. Add the remaining seed to the planter box and treat by thoroughly stirring with a paddle or lath so that the seed is coated with the seed treatment.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.