

Hombre Pellets

Ready-to-Use

Kills Norway Rats, Roof Rats, House Mice and Warfarin-Resistant House Mice
{Rats and Mice may consume a lethal dose in a single feeding}

COMMERCIAL

GUARANTEE: Difethialone..... 0.0025%

REGISTRATION NO. 26575 PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING
WARNING - CONTAINS THE ALLERGENS PEANUTS AND WHEAT

WARNING
[Diamond Skull & Crossbones Symbol]
POISON

Liphatech Inc.
3600 W. Elm Street
Milwaukee, WI
53209 U.S.A.
(414) 351-1476

NET CONTENTS: 15 kg

DIRECTIONS FOR USE:

For the control of Norway rats, Roof rats and House mice inside and in areas adjacent to homes, industrial, commercial, agricultural and public buildings.

Selection of Treatment Areas: Before placing bait, eliminate as much of the naturally occurring food as possible. Place bait in locations where rats and mice travel and will most likely find and consume the bait. Place bait only inside the building and in areas adjacent to buildings. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. For roof rats, bait placement should also be made in areas other than at ground/floor level. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

Treated baits should be placed in tamper-resistant bait stations, or in locations not accessible to children, pets, wildlife or domestic animals. Tamper-resistant bait stations must have the following characteristics: be constructed of high-strength material and resistant to destruction by children and animals; have an entrance which prevents children and non-target animals from reaching bait; have an internal structure which prevents bait from being shaken loose; have an access panel which fastens securely and locks; capable of being securely fastened to a surface; be clearly labelled: DANGER POISON. Note: This product is not to be used in orchards, fields or rights-of way. Use in conjunction with rat-proofing methods if rodent problems persist.

Food Processing, Food Manufacturing, Food Storage and Food Service Areas: For areas not directly related to food processing: Use only in non-food or non-feed areas where feed, food, packaging and handling equipment are never opened or exposed. For areas where feed or food is processed, served or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents and wash treated areas before re-use of the food handling establishment (processing areas include storage and service).

APPLICATION DIRECTIONS:

RATS: Place at least 85 to 170 grams of bait per placement at intervals of 5 to 10 meters. Larger amounts (up to 450 grams) may be necessary at points of very high rat activity. Maintain an uninterrupted supply of fresh bait for 10 days, or until signs of rat activity cease.

MICE: Place 7 to 14 grams of bait per placement at intervals of 2 to 4 meters. Larger amounts (up to 56 grams) may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days, or until signs of mouse activity cease.

Note: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate rodent food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS AND WILDLIFE. Keep away from feed or food. May irritate eyes. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. USE GLOVES WHEN HANDLING PRODUCT AND WHEN DISPOSING OF DEAD RODENTS, UNCONSUMED BAITS AND EMPTY CONTAINERS. Do not get in eyes, on skin, or on clothing. Wash arms, hands, and face with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the application occurs.

FIRST AID:

If in eyes, flush with water for 15 minutes. If irritation persists, seek medical attention. If on skin, wash affected area thoroughly with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object; repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Seek medical attention or contact Poison Control Centre IMMEDIATELY. Take container, label or product name and Pest Control Products Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments. Contains peanuts and wheat - may cause allergic reaction in sensitive persons.

STORAGE AND DISPOSAL:

Store product not in use in original, tightly sealed container in a cool, dry, secure location inaccessible to children, pets, domestic animals or wildlife. Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

RODENT DISPOSAL:

To reduce exposure of non-target birds and other wildlife to poisoned carcasses, dead rodents should be securely wrapped and placed in closed containers and disposed of at Provincial sites, where available, or at municipal landfill sites. Otherwise, securely wrap carcasses and bury to a depth that will make them inaccessible to scavengers.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.