

2016-3570
2017-01-31

MilStop®

Foliar Fungicide

MilStop Foliar Fungicide is a contact fungicide for control/suppression of Powdery Mildew on Greenhouse Vegetable Crops, Greenhouse Ornamental Crops, Outdoor Bedding Plants (including, but not limited to, African Daisy, Poinsettia, Hydrangea, Phlox, Rose, Bee Balm, Flowering Dogwood), Field Cucurbits (Cucumbers, Melons, Pumpkins, Squash), Stone Fruits (Peach, Nectarine, Apricot, Plum, Prune), Cannabis (marihuana) that is produced commercially indoors, Hops, Greenhouse and Field Strawberries, Greenhouse and Field Herbs and Spices; Grapes (conventional and organic), Artichokes, and Field Peppers.

COMMERCIAL

AL DUST

**WARNING SKIN
IRRITANT CAUTION
EYE IRRITANT
WARNING CONTAINS THE ALLERGEN**

SULFITES READ THE LABEL BEFORE USING

GUARANTEE : Potassium bicarbonate

85.00% REGISTRATION No. 28095 PEST

CONTROL PRODUCTS ACT

NET CONTENTS :

BioWorks, Inc. 100 Rawson Road, Suite 205, Victor, N.Y. 14564, Telephone 1-800-877-9443

GENERAL INFORMATION: MilStop Foliar Fungicide is a contact fungicide that controls/suppresses powdery mildew on greenhouse vegetable crops and ornamental crops; suppresses powdery mildew on outdoor bedding plants, field cucurbits stone fruits, Cannabis (marihuana) that is produced commercially indoors, hops, greenhouse and field strawberries, and greenhouse and field herbs and spices; and controls powdery mildew on grapes (conventional and organic), artichokes and field peppers. MilStop Foliar Fungicide is applied using a sufficient volume of water to insure complete coverage of all stems, foliage and fruit.

IMPORTANT : Read the entire label before using MilStop Foliar Fungicide.

DIRECTIONS FOR USE : Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. DO NOT allow effluent or runoff containing this product to enter lakes, streams, ponds or other waters. As this pesticide is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

DO NOT apply by air.

KEEP OUT OF REACH OF CHILDREN

FIRST AID:

INGESTION: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

INHALATION: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN/CLOTHING: Take off contaminated clothing. Rinse skin IMMEDIATELY with plenty of water for 15 – 20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for further treatment advice.

GENERAL: Remove victim to a safe and uncontaminated area. Seek medical attention or contact a poison control centre IMMEDIATELY if irritation occurs and persists or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. There is no need to induce vomiting. Monitor patient until ingested material clears their system.

PRECAUTIONS

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreation areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid contact with eyes. May irritate eyes. Causes skin and mucous membrane irritation. DO NOT get on skin. Avoid breathing dust or spray mist. Do not store near food or feed.

ENVIRONMENTAL HAZARDS: If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

USER SAFETY RECOMMENDATIONS: Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then thoroughly wash and put on clean clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear appropriate eyewear, waterproof gloves, long-sleeved shirt and pants or coveralls, and footwear with socks.

There is a 4-hour restrictive entry-interval (REI) requirement. Personal protective equipment is required for entry into treated areas that may involve contact with anything that has been treated, such as plants, soil or water. Individuals should wear gloves, long-sleeved shirt and pants or coveralls, and shoes plus socks for any post-application activities.

MIXING INSTRUCTIONS: MilStop Foliar Fungicide must be diluted with water before applying. Carefully measure and mix the recommended amounts of product and water. Fill tank with $\frac{1}{2}$ to $\frac{3}{4}$ of the selected volume of water prior to the addition of MilStop Foliar Fungicide. Start mechanical or hydraulic agitation. Slowly add the required amount of MilStop Foliar Fungicide to the tank. Continue to fill the tank with water to the desired volume while agitating to promote uniform distribution and prevent settling.

DO NOT MIX WITH OTHER SOLUBLE PESTICIDES OR FERTILIZERS NOT COMPATIBLE WITH MILD ALKALINE SOLUTIONS. THE ADDITION OF SURFACTANT OR WETTING AGENT TO SPRAY IS NOT REQUIRED. PRODUCT IS PRE-FORMULATED WITH THESE PRODUCTS FOR BEST PERFORMANCE. DO NOT ADJUST SPRAY SOLUTION pH AFTER MIXING OF PRODUCT. ACIDIFICATION OF SOLUTION WILL CAUSE REDUCED PRODUCT PERFORMANCE.

SPRAY APPLICATION METHOD: Do not apply this product through any type of irrigation system. For best results, use a properly calibrated sprayer and check the calibration frequently.

AGRICULTURAL APPLICATIONS: MilStop Foliar Fungicide is recommended for the control/suppression of Powdery mildew caused by: *Oidium lycopersicum* on tomato; *Leveillula taurica* on artichokes and sweet pepper; *Phodosphaera xanthii* (formerly named *Sphaerotheca fuliginea*) on cucumber and pumpkin grown in greenhouses; *Podosphaera clandestina* and *Sphaerotheca pannosa* on stone fruits (peach, nectarine, apricot, plum, prune); *Podosphaera spp.* and *Erysiphe spp.* on field cucurbits; *Uncinula necator* on grapes (conventional and organic); *Podosphaera macularis* (formerly *S. humuli*) on hops; *Sphaerotheca spp.*, *Erysiphe Spp.*, *Podosphaera spp.* on Greenhouse and Field Strawberries and Herbs and Spices; *Sphaerotheca spp.* on Cannabis (marihuana) that is produced commercially indoors.

ORNAMENTAL APPLICATIONS: MilStop Foliar Fungicide is recommended for the control/suppression of Powdery mildew caused by: *Oidium* sp. On African daisy and poinsettia; *Erysiphe polygoni* on hydrangea; *Erysiphe cichoracearum* on phlox and bee balm; *Microsphaera* sp. on flowering dogwood; and for the suppression of Powdery mildew caused by: *Oidium* sp., *Erysiphe polygoni*, *Erysiphe cichoracearum*, *Microsphaera* sp. and *Sphaerotheca pannosa* on outdoor bedding plants.

GREENHOUSE APPLICATION: Apply at a rate of 0.28 to 0.56 kilograms per 1000 sq. m. or 1/10 hectare. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Number of applications will depend upon disease pressure. Do not exceed a use rate of 0.56 kilograms of MilStop Foliar Fungicide per 1000 sq. m. or 1/10 hectare, per application. DO NOT STORE UNUSED SPRAY SOLUTION. DISPOSE OF UNUSED SOLUTION ON SITE.

FIELD/OUTDOOR APPLICATION: Apply at a rate of 2.8 to 5.6 kilograms of MilStop Foliar Fungicide per hectare. Use of the lower rate (2.8 kg/ha) may reduce damage from powdery mildew infections in stone fruits. Use a minimum water volume of 1000 L/ha to obtain complete coverage of foliage, stems and fruit. Uniform and complete coverage of the foliage, stems and fruit are essential for the most effective results. Spray volume per hectare can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 5.6 kilograms of MilStop Foliar Fungicide per hectare per application. DO NOT STORE UNUSED SPRAY SOLUTION. DISPOSE OF UNUSED SOLUTION ON SITE.

MAXIMUM NUMBER OF APPLICATIONS PER SEASON: Ten (10)

Not all bedding plants or varieties of bedding plants have been tested for phytotoxicity. It is recommended that a phytotoxicity test be conducted on a small group of representative plants before applying to a large number of plants. MilStop Foliar Fungicide may cause phytotoxicity on certain cultivars of hydrangea.

APPLICATION TIMING: Start application at first sign of disease. For best protection, repeat at one or two week intervals as indicated until conditions are no longer favorable for disease development. Use the high rate (5.6 kg/ha) under conditions of high to moderate disease pressure and the low rate (2.8 kg/ha) under conditions of low to moderate disease pressure. Shorten the application interval to 7 days during rainy weather or during periods of high relative humidity. The pre-harvest interval (PHI) is 0 days.

Crops	Use sites	Disease	Pathogens	Rate (kg/ha)	Days between applications	Spray volume (L/ha)	Control/suppression
<u>Vegetables</u>	Greenhouse	Powdery mildew					
Cucumber, Sweet Pepper, Tomato			<i>Podosphaera xanthii</i> , <i>Leveillula taurica</i> , <i>Oidium lycopersicon</i>	5.6	7	2000	Control
Pumpkin			<i>Podosphaera xanthi</i>	5.6	7	1000	Suppression

Ornamentals	Greenhouse	Powdery Mildew						
Flowering Dogwood, Bee Balm			<i>Microsphaera</i> spp., <i>Erysiphe cichoracearum</i>	5.6	7 - 14	1000	Control	
African Daisy			<i>Oidium</i> spp.	2.8 - 5.6	7 - 14	1000	Control	
Hydrangea, Poinsettia,	Field	Powdery Mildew	<i>Erysiphe polygoni</i> , <i>Oidium</i> spp.	5.6	7 - 14	2000	Control	
Phlox			<i>Erysiphe cichoracearum</i>	2.8 - 5.6	7 - 14	1000	Suppression	
Bedding Plants Including but not limited to:								
Flowering Dogwood, Bee Balm				<i>Microsphaera</i> sp., <i>Erysiphe cichoracearum</i>	2.8 - 5.6	7 - 14	1000	Suppression
African daisy				<i>Oidium</i> sp.	2.8 - 5.6	7 - 14	1000	Suppression
Hydrangea, Poinsettia				<i>Erysiphe polygoni</i> , <i>Oidium</i> sp.	2.8 - 5.6	7 - 14	1000	Suppression
Phlox				<i>Erysiphe cichoracearum</i>	2.8 - 5.6	7 - 14	1000	Suppression
Rose		<i>Sphaerotheca pannosa</i>	2.8 - 5.6	7 - 14	1000	Suppression		
Grapes	Field	Powdery mildew	<i>Uncinula necator</i>	2.8 - 5.6	7 - 14	500 - 1000	Control	
Cucurbits	Field	Powdery mildew	<i>Podosphaera</i> spp., <i>Erysiphe</i> spp.	2.8 - 5.6	7 - 14	500 - 1000	Suppression	
Artichokes	Field	Powdery mildew	<i>Leveillula taurica</i> , <i>Erysiphe cichoracearum</i>	2.8 - 5.6	7 - 14	1000	Control	
Field Peppers	Field	Powdery mildew	<i>Leveillula taurica</i>	2.8 - 5.6	7	1000	Control	
Stone Fruits	Field	Powdery mildew	<i>Podosphaera clandestine</i> , <i>Sphaerotheca pannosa</i>	2.8 - 5.6	7 - 14	1000	Suppression	
Hops	Field	Powdery mildew	<i>Podosphaera macularis</i>	2.8 - 5.6	7 - 14	500 - 1000	Suppression	

Cannabis (marihuana)	This product is to be used only on cannabis (marihuana) that is produced commercially indoors.	Powdery mildew	<i>Sphaerotheca macularis.</i>	0.28-0.56 kg/1000 m ²	7 – 14	1000	Suppression
Strawberries	Greenhouse or Field	Powdery mildew	<i>Sphaerotheca macularis</i>	2.8 – 5.6	7 - 14	1000	Suppression
Herbs and Spices*	Greenhouse or Field	Powdery mildew**	<i>Sphaerotheca spp., Erysiphe spp., Podosphaera spp.</i>	Greenhouse: 0.28-0.56kg/1000m ² Field:2.8 - 5.6	7 - 14	1000	Suppression

*angelica, anise, balm, basil, borage, burnet, chamomile, caraway, black caraway, catnip, celery seed, chervil, chive, Chinese chive, clary, coriander, costmary, cilantro, cumin, dill, fennel, Florence fennel, fenugreek, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, marigold, marjoram, mustard, nasturtium, parsley, pennyroyal, poppy, rosemary, rue, saffron, sage, savory (summer, winter), sweet bay (bayleaf), tansy, tarragon, thyme, woodruff, wormwood

**Herbs and herb varieties may vary in their tolerance to pest control products, including MILSTOP fungicide. Since not all herbs and herb varieties have been tested for tolerance to MILSTOP fungicide, first use of MILSTOP fungicide should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of herbs and herb varieties to MILSTOP fungicide.

STORAGE :

To prevent contamination, store this product away from food or feed.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**In Case of Emergency,
Call CHEMTREC: (800) 424-9300**

Manufactured in the USA by:

BioWorks, Inc.
100 Rawson Road
Suite 205
Victor, NY 14564
Telephone 1-800-877-9443